

Designing Psychologically Responsive Healthcare Systems: Integrating Behavioral Science, Equity, Emotional Needs, and Human-Centered Innovation

Luca Ferraro^{1*}, Matteo Ricci¹, Giulia Moretti², Paolo Greco³

¹Department of Health Systems Design and Behavioral Science, Faculty of Medicine, University of Bologna, Bologna, Italy.

²Department of Equity and Human-Centered Innovation, Faculty of Medicine, University of Turin, Turin, Italy.

³Department of Psychologically Responsive Care, Faculty of Medicine, Sapienza University of Rome, Rome, Italy.

*E-mail ✉ luca.ferraro@unibo.it

Abstract

Healthcare systems are commonly evaluated through access, safety, effectiveness, efficiency, and patient-centredness, yet these domains do not fully capture whether care environments can recognize and respond to the psychological conditions that shape participation in care. Emotional needs, trust, perceived legitimacy, health literacy, social resources, organizational culture, digital accessibility, and prior experiences influence how patients interpret options and whether opportunities to participate are practically usable. This conceptual article develops the idea of a psychologically responsive healthcare system as a system-level capacity rather than an individual communication technique or patient attribute. It integrates behavioral science, relational care, equity, human-centered service design, workforce conditions, digital health, and learning-system principles while maintaining distinctions among patient experience, psychological process, observable behavior, care-process outcomes, and clinical outcomes. The proposed account treats engagement capacity as relational and context-dependent, recognizes organizational conditions as constraints on frontline responsiveness, and positions digital technologies as sociotechnical environments rather than neutral delivery channels. It further argues that psychological responsiveness requires iterative learning because patient needs, resources, illness states, technologies, and care pathways change over time. The resulting framework is explicitly theoretical: available evidence supports many constituent relationships, but does not establish that their integration into a single architecture will necessarily improve clinical outcomes. The model therefore provides design principles, boundaries, and testable domains for future implementation and evaluation rather than claiming validated system effectiveness.

Keywords: Psychological responsiveness, Patient engagement, Relational care, Behavioral science, Health equity, Human-centered healthcare

Introduction

Patient-centredness has evolved into a multidimensional construct encompassing recognition of the patient as a person, attention to biopsychosocial context, therapeutic relationships, information exchange, and some redistribution of decision influence rather than a single

measure of satisfaction or communication quality [1]. Yet even a strongly patient-centred service may remain psychologically unresponsive when patients are technically offered choices but cannot understand, trust, access, or act on them. Shared decision-making illustrates this distinction: it is an important participatory process, but its effects are not uniformly positive across every health, quality, utilization, or cost outcome [2]. Consequently, participation itself should not be treated as proof that care has accommodated the psychological and contextual conditions required for meaningful engagement.

Patient involvement in service design and quality improvement can influence healthcare processes,

Access this article online

<https://smerpub.com/>

Received: 12 April 2026; Accepted: 01 August 2026

Copyright CC BY-NC-SA 4.0

How to cite this article: Ferraro L, Ricci M, Moretti G, Greco P. Designing Psychologically Responsive Healthcare Systems: Integrating Behavioral Science, Equity, Emotional Needs, and Human-Centered Innovation. *Int J Soc Psychol Asp Healthc*. 2026;6(2):98-112. <https://doi.org/10.51847/Y6K4VOYzeE>

although strategies, depth of involvement, and evaluated outcomes vary substantially [3]. Evaluation methods for patient and public engagement are likewise heterogeneous, making the existence of an engagement structure easier to document than its quality, influence, or distribution of power [4]. These limitations suggest a wider design problem. Healthcare systems must not only create opportunities for patient participation; they must also make those opportunities interpretable, relationally safe, practically accessible, and responsive to changing circumstances.

This article therefore proposes psychological responsiveness as a system property: the capacity to detect patient-relevant psychological and social conditions, interpret them without reducing them to fixed traits, adapt care processes within legitimate clinical boundaries, and learn when those adaptations succeed or fail. The concept does not imply that health systems can or should eliminate distress, guarantee trust, or individualize every decision. Instead, it directs attention to the fit between patients, relationships, services, technologies, organizations, and structural conditions. The argument integrates behavioral science, emotional and relational care, equity, human-centered design, workforce conditions, digital responsiveness, and continuous learning while separating established evidence from the new theoretical architecture developed here.

What makes a healthcare system psychologically responsive?

Psychological responsiveness begins where conventional patient-centredness becomes operationally demanding. A system must be able to recognize that two patients receiving the same clinical option may face different interpretive, emotional, social, linguistic, or material conditions for using it. Responsiveness therefore concerns more than whether information was delivered or a preference was recorded. It concerns whether the care environment makes meaningful participation possible and whether the system can adjust when its assumptions about patients prove inaccurate.

Formal structures alone are insufficient. Evaluation of patient participation in value-based healthcare has shown that participation can vary substantially even where organizations explicitly endorse patient involvement [5]. The implication is not that formal participation mechanisms are unimportant, but that their presence is an inadequate proxy for influence. A psychologically

responsive system must distinguish nominal opportunity from usable opportunity and attendance from consequential participation.

Responsiveness also extends beyond the conventional patient-professional dyad. Peer involvement can connect experiential knowledge, community relationships, and formal care, particularly where conventional institutions have weak reach or trust [6]. Such roles should not be romanticized: their functions vary across contexts, and peers cannot compensate indefinitely for inaccessible or discriminatory systems. Nevertheless, they demonstrate that psychologically meaningful information can enter care through social relationships that sit partly outside professional structures.

We therefore propose four linked capabilities. First, the system must sense patient-relevant needs and constraints without assuming that demographic identity reliably predicts psychological state. Second, it must interpret those signals in clinical and social context. Third, it must possess sufficient response capacity—relational, organizational, informational, and material—to modify care where appropriate. Fourth, it must revise its practices when patient feedback, outcome discordance, or changing circumstances reveal that the original response was inadequate. These capabilities are proposed analytical functions, not validated dimensions of a measurement scale.

Behavioral science as a design foundation

Healthcare frequently frames nonparticipation as deficient motivation: the patient did not adhere, engage, attend, disclose, or follow advice. Behavioral science complicates that interpretation. Applications of the Theoretical Domains Framework and related behavior-change approaches organize behavior across multiple determinants rather than treating motivation as the dominant explanation [7, 8]. Capability, knowledge, beliefs, professional or social roles, environmental resources, reinforcement, emotion, and opportunity may interact differently across settings.

For psychologically responsive design, the important implication is diagnostic. Before attempting to increase engagement, the system should ask what makes the target behavior feasible, meaningful, and safe. A missed appointment may reflect competing responsibilities or transportation constraints; apparent indecision may reflect uncertainty or previous experiences of dismissal; silence may represent fear of judgment rather than lack of information. These interpretations are hypotheses that

require contextual evidence, not assumptions to be assigned to groups.

Health literacy provides a clear example. Interventions intended to improve health literacy and chronic-disease self-management show that the ability to act on health information can be modified, but effects depend on how information and support are designed [9]. Thus, demands created by healthcare environments belong inside behavioral analysis. A service that requires high literacy, rapid decisions, digital navigation, and repeated self-advocacy may create behavioral burdens that are subsequently mislabelled as patient disengagement.

The design principle proposed here is therefore behavioral fit: services should diagnose the conditions governing participation before selecting behavioral interventions. Psychological responsiveness does not mean engineering compliance. It means reducing avoidable mismatch between what care requires and what patients can realistically understand, tolerate, access, and enact while preserving informed refusal and legitimate clinical constraints.

Emotional needs and relational care

Emotional responsiveness is often reduced to staff warmth, yet relational care involves more specific processes. Patient-informed work on compassionate healthcare indicates that compassion can be operationalized through observable actions rather than treated solely as a professional intention [10]. Trust is similarly multidimensional: it emerges from interacting expectations and experiences within the patient-physician relationship rather than from a single

communicative behavior [11]. A system therefore cannot assume that technically correct information, delivered politely, will automatically be experienced as trustworthy or supportive.

Empathy provides both supportive evidence and an important boundary. Randomized evidence indicates that practitioner-empathy interventions can improve patient satisfaction [12]. Satisfaction, however, is not equivalent to clinical outcome, durable trust, adherence, or system equity. Psychological responsiveness requires these outcomes to remain analytically separate. A patient may report a respectful encounter while still facing inaccessible follow-up care, or may distrust a recommendation despite excellent technical care because previous encounters have shaped expectations.

Compassion education further illustrates the difference between professional capability and system capacity. Training approaches can strengthen compassionate practice, but the evidence is heterogeneous and cannot establish that education alone overcomes workload, staffing, continuity, organizational, or resource constraints [13]. Relational quality is therefore partly produced by organizational conditions.

A responsive design should make emotional needs discussable without requiring emotional disclosure as a condition of receiving good care. It should allow uncertainty, fear, embarrassment, grief, loss of control, and concerns about legitimacy to be acknowledged when clinically relevant, while avoiding the assumption that every patient wants intensive emotional engagement. **Table 1** distinguishes these relational conditions from the contexts that change their meaning.

Table 1. Relational conditions that alter whether emotional responsiveness becomes usable care

Patient-facing process that matters	Context that can change its meaning	Competing interpretation or uncertainty	Boundary case	Design consequence
Relational condition — Feeling recognized as a person				
Attention to concerns, biography and expressed priorities	Time pressure, repeated handoffs, unfamiliar teams	Courtesy can be mistaken for genuine recognition	Some patients prefer narrowly task-focused encounters	Preserve opportunities for recognition without requiring personal disclosure
Relational condition — Trust under clinical uncertainty				
Transparency about reasoning, limits and next steps	Previous dismissal, diagnostic uncertainty, institutional reputation	Agreement does not necessarily indicate trust	Trust may remain low despite appropriate care	Make uncertainty and reasoning discussable rather than expecting automatic confidence
Relational condition — Experienced empathy				
Clinician responses that demonstrate understanding	Severity, emotional intensity, cultural	Patient satisfaction cannot be assumed to represent better clinical outcome	Neutral professional style	Evaluate relational experience separately

	expectations, continuity		may be preferred by some patients	from downstream outcomes
Relational condition — Compassion under operational pressure				
Timely supportive action, not intention alone	Workload, staffing, competing safety demands	Training effects may be limited by organizational constraints	Acute emergencies can legitimately prioritize rapid task performance	Treat compassionate practice as a workforce-and-system capability
Relational condition — Emotional safety to ask or disagree				
Nonpunitive questioning, clarification and dissent	Power asymmetry, stigma, language discordance, prior discrimination	Silence may reflect many causes and cannot be psychologized automatically	Patients may choose not to participate in deliberation	Create psychologically safer options for participation while preserving refusal
Relational condition — Proposed responsive condition				
Relational response matched to expressed need	Patient preference and changing illness trajectory	Responsiveness cannot be inferred from one interaction	Needs can change across encounters	Reassess relational fit rather than fixing patients into stable preference categories

Equity and differential capacity to engage

Engagement is often described as though every patient receives the same opportunity and differs mainly in willingness to use it. Equity evidence challenges that assumption. Patients' accounts of diversity-sensitive care emphasize that respectful communication, recognition of individual circumstances, and adaptation to differing needs matter, while demographic categorization alone cannot specify what an individual patient requires [14]. Responsiveness must therefore accommodate difference without converting population-level patterns into stereotypes.

Language is one concrete system condition. Evidence linking language discordance with unplanned readmissions or emergency revisits indicates that communication barriers can extend beyond subjective experience into subsequent care processes, although the underlying evidence is primarily observational and context dependent [15]. Likewise, qualitative synthesis in mental healthcare demonstrates how intersecting forms of discrimination can operate through both interpersonal encounters and structural arrangements [16]. These findings support multilevel attention to power without implying that every member of a marginalized group has the same experience.

Accessibility also concerns how information is represented. Visual approaches can improve aspects of health comprehension compared with some conventional communication formats, with effects dependent on the comparator and context [17]. Digital-health research similarly identifies interacting cultural, educational,

residential, socioeconomic, health, and technological conditions shaping use [18]. Consequently, “capacity to engage” is better treated as an emergent relation between person and system than as an intrinsic patient characteristic.

The proposed equity principle is differential usability. A service is not equitable merely because the same pathway is offered to everyone. Psychological responsiveness requires asking who can interpret the pathway, who bears its administrative and emotional burdens, whose preferences are recognized as legitimate, and who has realistic alternatives when the default pathway fails.

Human-centered service design

Human-centered design offers methods for translating patient experience into service architecture, but participation alone does not establish responsiveness. Evaluation research on health co-design emphasizes the need to assess both the co-design process and its effects rather than assuming benefit from involvement itself [19]. In multimorbidity, co-design has produced promising improvements in relevance, usability, and acceptability, while evidence of direct health benefit remains limited [20]. This distinction is central: better design experience is valuable but should not be relabelled as proven clinical effectiveness.

Power also shapes co-design. Health-policy co-design can create meaningful participation, yet implementation strategies differ in how far decision influence is transferred to patients and communities [21]. Applied human-centered work with patient advisory groups

shows the practical importance of iteration, relationship building, and consensus while also revealing time and coordination constraints [22]. Digital co-design adds further challenges involving recruitment, representativeness, usability testing, technological feasibility, and scaling [23].

A psychologically responsive design process must therefore connect four questions: whose experience is sensed, whose interpretation shapes the problem

definition, whose constraints influence the prototype, and whose outcomes determine whether the design is retained? The answer cannot be “the patient” in the abstract. Different people may have conflicting needs, and clinical evidence, legal duties, workforce constraints, accessibility, and resource stewardship remain legitimate design boundaries.

Figure 1 maps these boundaries as nested contexts rather than a linear co-design sequence.




Figure 1. Participation Without Influence: Boundaries Separating Human-Centered Design From Psychologically Responsive Service Design

Organizational culture and workforce conditions

Psychological responsiveness is often expected from individual clinicians while the organization determines how much time, continuity, discretion, and emotional capacity they possess. Physician burnout is associated with patient-safety outcomes, professionalism, and patient satisfaction, but much of this evidence is observational and should not be interpreted as proving a simple causal pathway from clinician distress to every patient outcome [24]. Its stronger implication for this

framework is that relational performance cannot be detached from working conditions.

Organizational culture also matters. Leadership, teamwork, communication, recognition, autonomy, and broader workplace conditions are associated with provider work satisfaction [25]. Yet changing organizations is not straightforward. Systematic evidence on organization-directed well-being interventions reports heterogeneous results and generally low certainty, including neutral and occasionally unfavorable findings [26]. Reviews focused on primary care likewise support

addressing workload and organizational determinants rather than relying solely on individual coping strategies, while transferability remains context specific [27]. Psychological responsiveness should therefore not become another demand imposed on exhausted staff. We propose response capacity as the organizational resources that allow clinicians and teams to notice concerns,

communicate uncertainty, adapt processes, coordinate care, and repair relational ruptures without compromising safety. The concept includes time and staffing but is not reducible to either. **Table 2** uses an exception-first structure to show how apparent interpersonal failures may originate elsewhere in the organization.

Table 2. When Relational Failure Is Organizational: Exception-First Conditions That Constrain Psychological Responsiveness

Failure first observed in care	Latent organizational condition	Patient-facing manifestation	Misleading explanation to avoid	Exception or uncertainty	Responsive redesign priority
Patient concerns repeatedly re-explained	Fragmented continuity or handoffs	Fatigue, reduced disclosure, perceived nonrecognition	“Patient is difficult or repetitive”	Handoffs may be clinically necessary	Improve information continuity and ownership
Clinician appears rushed or emotionally unavailable	Excess workload or competing task demands	Reduced opportunity for questions and relational repair	“Clinician lacks empathy”	Brief encounters can still be respectful and effective	Protect high-value relational time where feasible
Staff training produces little visible change	Workflow and incentive structures remain unchanged	Compassionate intentions are inconsistently enacted	“Training failed because staff resisted”	Training content or delivery may also be inadequate	Align work design with expected relational behaviors
Patient receives inconsistent explanations	Weak team communication or role clarity	Uncertainty and reduced confidence	“Patient misunderstood”	Clinical uncertainty can legitimately change recommendations	Make uncertainty and responsibility for updates explicit
Well-being intervention has neutral or adverse effects	Poor fit between intervention and local job demands	No increase in staff response capacity	“More resilience training is required”	Effects differ across settings and intervention designs	Diagnose modifiable organizational demands before adding interventions
Proposed system-level failure mode	Responsiveness expectations exceed available capacity	Relational quality becomes episodic and dependent on individual sacrifice	“Good clinicians will compensate”	Some high-performing teams may temporarily buffer constraints	Evaluate system capacity instead of normalizing compensatory overwork

Digital and technological responsiveness

Digital healthcare can expand contact while simultaneously relocating work and risk to patients. Rapid telehealth expansion demonstrated that access depends on modality, infrastructure, workflow, and local social conditions rather than on service availability alone [28]. Health literacy and area deprivation have also been associated with whether telehealth visits are initiated and completed, reinforcing the importance of person-system

fit [29]. Digital engagement should therefore not be interpreted as a simple preference for or rejection of technology.

The same principle applies to professionals. Adoption of digital technologies is influenced by infrastructure, workload, training, usability, organizational support, and perceived value [30]. Digital-health-equity frameworks consequently operate across multiple levels rather than through a single “access” construct [31]. Trust introduces

another layer: consumer and professional trust in digital healthcare encompasses privacy, reliability, accuracy, human interaction, literacy, and perceptions of system performance [32]. Current digital-equity frameworks and inequality research together indicate that narrow attention to access or use can miss broader distributional consequences [31, 33].

Importantly, disadvantage does not produce uniform patterns of digital nonuse. Recent population evidence shows heterogeneous associations between individual characteristics, place-level vulnerability, and use of different digital-health modalities [34]. Demographic variables should therefore guide investigation of potential inequities, not determine assumptions about individual capability or preference.

Psychologically responsive digital design should provide intelligible choices among modalities, preserve human escalation when automated or remote communication becomes insufficient, and assess the burdens transferred to patients. Evidence standards should also match the maturity and risk of the technology rather than treating adoption as evidence of effectiveness [35]. Digital responsiveness is thus a sociotechnical property: technology, patient capability, relational trust, professional workflow, accessibility, and organizational governance must be evaluated together while remaining analytically distinct.

Learning, feedback, and adaptive improvement

A psychologically responsive service cannot be fully specified once and then implemented unchanged. Patient needs evolve with illness, treatment, family circumstances, digital capability, accumulated care experiences, and institutional change. Learning-health-system scholarship describes an iterative relationship between data, knowledge generation, implementation, and improvement, while implementation-focused synthesis emphasizes the infrastructure needed for adaptation, scale-up, and sustainment [36, 37].

For psychological responsiveness, however, the content entering the learning loop matters as much as the existence of the loop. Clinical outcomes and utilization data may reveal whether care was effective or used, but they may not explain why a pathway was abandoned, why trust deteriorated, why a technically available service remained inaccessible, or why a change improved experience for one group while increasing burden for another. Patient-reported experience, qualitative feedback, complaints, accessibility failures, workforce observations, and distributional patterns therefore provide complementary—not interchangeable—signals. The framework proposed here treats feedback as a mechanism for recalibration. A responsive organization should be able to distinguish a one-off mismatch from a recurring design failure, determine the level at which correction is required, and revisit earlier interpretations when downstream experience contradicts them. **Table 3** makes this temporal logic explicit.

Table 3. Recalibration Across Time: How Psychological-Responsiveness Signals Should Change System Decisions

Transition point	Information that must remain visible	Failure produced by delay or omission	Adaptive action	Equity safeguard	Evidence status
Initial service encounter → early use	Expressed concerns, comprehension, preferred participation, practical access	Early mismatch becomes interpreted as nonadherence	Clarify demands and adjust communication or pathway where appropriate	Do not infer capability from demographic category	Evidence-supported premises; transition logic proposed
Early use → continuing care	Experience of trust, burden, continuity and feasibility	Repeated friction accumulates without formal recognition	Reassess relational and logistical fit	Identify who bears extra administrative or digital work	Proposed integration
Clinical or life-state change → renewed decision	Changed illness state, emotional priorities, social resources, treatment demands	Previously recorded preferences become treated as permanent	Re-open deliberation and update support needs	Preserve accessible routes back into decision-making	Proposed temporal principle

Service redesign → implementation	Usability, staff workflow, unintended exclusions, clinical constraints	Design success is inferred from launch or uptake	Compare intended design with implementation experience	Examine differential reach and burden	Supported implementation principle
Implementation → organizational learning	Patient feedback, workforce signals, complaints, outcomes and distributional patterns	Data accumulate without changing practice	Route discordant signals to the level able to act	Avoid averaging away subgroup or accessibility failures	Supported learning principle with proposed psychological extension
Apparent success → periodic reassessment	Benefits, trade-offs, excluded populations, changing technology/context	Temporary fit is mistaken for durable responsiveness	Revalidate assumptions and retire ineffective adaptations	Reassess who is no longer served by the prevailing design	Proposed recalibration rule

A learning architecture of this kind does not guarantee improvement. Feedback can be selective, data can reproduce existing exclusions, and organizations may collect information without possessing the authority or resources to respond. Psychological responsiveness therefore requires not simply more measurement, but the capacity to connect discrepant signals with accountable decisions and to recognize when the relevant problem lies at a different analytical level from the one first observed.

Integrating clinical, behavioral, social, and organizational levels

Psychological responsiveness fails when phenomena observed at one level are automatically explained at the same level. A patient's missed follow-up can appear behavioral while arising from inaccessible scheduling, unstable transport, caring responsibilities, digital demands, or a previous relational rupture. Conversely, structural disadvantage does not determine an individual's beliefs, emotions, or preferences. Multilevel explanation therefore requires connection without collapse.

Experiential knowledge illustrates this principle. Models integrating patients' lived expertise with professional care show that experiential knowledge can contribute to chronic-care relationships without becoming interchangeable with clinical expertise [38]. Similar distinctions apply throughout the proposed framework. Behavioral science helps identify conditions governing action; relational evidence clarifies trust, empathy, communication, and recognition; organizational evidence identifies workforce and workflow constraints; equity analysis locates unequal burdens and power;

digital evidence reveals sociotechnical conditions; and clinical expertise defines legitimate diagnostic, therapeutic, and safety boundaries.

The proposed integration therefore uses cross-level correspondence rather than a hierarchy with one master cause. A psychological signal may require a relational response, an organizational redesign, or both. Persistent misunderstanding across many patients may indicate an information-system problem rather than repeated individual deficits. Recurrent clinician inability to provide adequate discussion time may indicate workload design rather than insufficient communication training. The analytical task is to identify the level at which a problem becomes visible, distinguish that from the level at which it is generated, and locate the level with authority to change it.

The psychologically responsive healthcare system framework

The evidence assembled above supports constituent principles but does not establish a validated unified system model. We therefore propose the Psychologically Responsive Healthcare System Framework as an integrative architecture with five interacting functions.

The first is sensing: recognizing patient-relevant emotional, behavioral, informational, relational, accessibility, and social signals while avoiding demographic stereotyping. The second is interpretation: determining what those signals may mean in the current illness, decision, relationship, and care context. The third is response capacity: possessing sufficient relational skill, clinical flexibility, organizational resources, accessible communication, and technological

alternatives to act when adaptation is warranted. The fourth is consequential participation: ensuring that patient or community input can influence decisions at the level where participation has been invited rather than being collected symbolically. The fifth is recalibration: using subsequent experience, outcomes, complaints, workforce information, accessibility failures, and distributional patterns to revise previous assumptions. These functions interact but should not be represented as a universal sequence. A system may sense a problem but lack authority to respond. Participation may occur before adequate interpretation. A digital redesign may improve accessibility for one population while introducing new burdens elsewhere. A clinically appropriate decision may remain emotionally difficult despite excellent relational care. Responsiveness therefore denotes the ability to accommodate relevant psychological and contextual conditions **within** clinical, ethical, organizational, and

resource boundaries, not an obligation to satisfy every preference.

The framework also distinguishes two processes that can either converge or diverge. One process concerns the patient's usable capacity to participate, shaped by comprehension, emotional state, trust, social resources, accessibility, illness burden, and previous experience. The other concerns the system's capacity to respond, shaped by clinical flexibility, workforce conditions, continuity, communication infrastructure, digital design, organizational authority, and learning capability. Psychological responsiveness is most plausible where these processes achieve sufficient fit. Divergence occurs when participation is demanded without support, when the system offers adaptations that patients cannot use, or when patient needs are recognized but organizational constraints prevent action.

Table 4 identifies where constraints arising at different levels can interrupt this fit.

Table 4. Multilevel Constraints That Can Break the Fit between Patient Participation and System Response

Participation signal	Patient capability	System response	Organization constraint	Mismatch effect	Recalibration priority
Level where constraint originates — Individual and illness context					
Condition visible to the patient or team	Pain, fatigue, fear, cognitive load, changing priorities	What becomes restricted	Ability or desire to process information and participate	Level that may need to respond	Encounter and pathway design
Interpretation boundary	Reduced participation must not automatically be labelled unwillingness	Proposed framework implication	Reassess participation demands as states change		
Level where constraint originates — Relational encounter					
Condition visible to the patient or team	Low trust, anticipated judgment, weak continuity, unresolved disagreement	What becomes restricted	Disclosure, questioning and collaborative deliberation	Level that may need to respond	Clinician/team relationship
Interpretation boundary	Agreement is not proof of trust; silence is not proof of understanding	Proposed framework implication	Create routes for clarification, dissent and relational repair		
Level where constraint originates — Social and material context					
Condition visible to the patient or team	Caring duties, transport difficulty, unstable resources, limited support	What becomes restricted	Practical ability to enact agreed care	Level that may need to respond	Service organization and cross-sector support
Interpretation boundary	Behavioral noncompletion does not establish motivational deficit	Proposed framework implication	Distinguish action capability from stated intention		
Level where constraint originates — Organizational environment					

Condition visible to the patient or team	Workload, fragmented handoffs, rigid workflows, limited discretion	What becomes restricted	Staff capacity to adapt care	Level that may need to respond	Team, management and governance
Interpretation boundary	Repeated interpersonal failure may have organizational origins	Proposed framework implication	Evaluate response capacity rather than relying on individual compensation		
Level where constraint originates — Digital environment					
Condition visible to the patient or team	Inaccessible interfaces, modality mismatch, privacy concern, high literacy demand	What becomes restricted	Entry to care, communication and follow-up	Level that may need to respond	Technology, workflow and service-design functions
Interpretation boundary	Digital availability is not equivalent to usable access	Proposed framework implication	Maintain alternative routes and monitor transferred burden		
Level where constraint originates — Structural and equity context					
Condition visible to the patient or team	Language barriers, discrimination, unequal influence, resource asymmetry	What becomes restricted	Recognition, legitimacy and consequential participation	Level that may need to respond	Organization, system and policy
Interpretation boundary	Population-level inequality cannot specify an individual's psychological state	Proposed framework implication	Examine distribution of burdens, voice and benefit		
Level where constraint originates — Learning system					
Condition visible to the patient or team	Feedback not captured, averaged away, delayed or disconnected from authority	What becomes restricted	Correction of recurring mismatches	Level that may need to respond	Quality, governance and implementation infrastructure
Interpretation boundary	Measurement without action is not responsiveness	Proposed framework implication	Link discrepant signals to accountable revision		

Figure 2 represents the proposed framework as paired fields whose convergence depends on cross-level conditions rather than as a deterministic causal pathway.




Figure 2. Where Patient Participation Capacity and Healthcare Response Capacity Converge—and Where They Separate

Design principles and evaluation domains

The framework implies that psychological responsiveness should be evaluated as a profile rather than a single score. Existing engagement measures already demonstrate that participation requires multidimensional evaluation [4], while patient-informed compassion measurement shows the value of specifying observable relational processes [10]. Communication accessibility should likewise be assessed separately from whether information was merely provided [17]. Five design principles follow. Design for usable participation: evaluate comprehension, practical feasibility, and acceptable alternatives rather than availability alone. Protect consequential voice: determine whether patient or community input can change decisions within the stated scope of participation [19]. Match demands to response capacity: avoid asking clinicians or patients to compensate indefinitely for poorly designed systems. Test distribution, not only averages: digital-equity frameworks and digital-trust evidence show why access, trust, burden, and benefit should be evaluated

across relevant populations without deterministic demographic assumptions [31]. Build for revision: evaluation should identify when changing circumstances make previous accommodations ineffective. Evaluation should accordingly distinguish at least relational experience, participation quality, behavioral feasibility, accessibility, workforce response capacity, digital trust and usability, organizational adaptability, and learning-system performance. Digital evidence standards further indicate that claims should be proportionate to evidence maturity and decision risk [35]. No composite index is proposed here because weighting these domains would introduce value judgments and empirical assumptions not established by the present evidence.

Research and implementation agenda

The immediate research requirement is not to test the entire framework as though it were a standardized intervention. Constituent propositions should first be operationalized and examined separately. Studies could

test whether mismatches between participation demands and available support predict disengagement more usefully than patient-level motivation alone, whether repeated relational problems cluster with measurable organizational constraints, and whether feedback systems detect distributional or accessibility failures before they become established pathway inequalities.

Longitudinal designs are particularly important because participation capacity and response capacity change. Co-design research demonstrates that process quality and clinical effectiveness cannot be assumed to coincide [20], while organization-directed workforce interventions show why plausible system changes can produce heterogeneous or null results [26]. Digital-health research similarly requires assessment across implementation maturity, setting, accessibility, and trust rather than simple adoption counts.

Implementation studies should specify which component is changing, which level possesses authority to act, what competing explanation remains plausible, and which outcomes are patient reported, behavioral, organizational, utilization related, or clinical. Comparative studies should examine whether the same design performs differently across cultural, linguistic, socioeconomic, disease, and service contexts. Equity evaluation should examine differential benefit and burden without treating group membership as an individual psychological diagnosis.

Ultimately, prospective testing should determine whether the proposed functions—sensing, interpretation, response capacity, consequential participation, and recalibration—are distinguishable, measurable, and useful. Failure to validate one component should prompt revision of the framework rather than post hoc relabelling of unsuccessful implementation as insufficient responsiveness.

Results and Discussion

The central argument of this article is that healthcare cannot be psychologically responsive through communication technique alone. Available evidence supports several constituent premises: patient-centredness is multidimensional [1]; meaningful participation differs from formal involvement [5]; behavioral engagement is shaped by more than motivation [7]; trust and compassion are relationally enacted [11]; equity conditions alter the usability of care [18]; co-design requires scrutiny of influence and

outcomes [19]; organizational conditions constrain frontline practice [25]; and digital participation is sociotechnical rather than purely technological [32]. The proposed framework integrates these premises but does not convert their coexistence into proof of a unified causal mechanism.

This distinction matters because several literatures contain important counterweights. Empathy can improve satisfaction without establishing broader clinical benefit [12]. Co-design can improve relevance or usability while evidence of health benefit remains limited [20]. Organizational interventions intended to improve workforce well-being can have heterogeneous, neutral, or unfavorable effects [26]. Digital inequalities cannot be predicted reliably from demographic disadvantage alone [34]. Psychological responsiveness should therefore not become a label applied retrospectively to every successful intervention.

A further risk is excessive personalization. Responsive care does not imply unlimited customization or the replacement of evidence-based care with preference satisfaction. Clinical safety, fairness, resource stewardship, professional duties, and population needs remain legitimate constraints. Nor should psychological explanation obscure structural causes. When a pathway is linguistically inaccessible or administratively burdensome, describing affected patients as anxious, disengaged, or low-literacy may individualize a system-generated obstacle.

The framework's practical value lies instead in improving diagnosis of mismatch. It asks whether a problem observed in patient behavior originates partly in relational, organizational, digital, or structural conditions; whether the system has capacity to respond; whether participation is consequential; and whether feedback can trigger revision. These questions are deliberately testable. Until prospective and comparative evaluation establishes otherwise, the framework should be treated as an organizing theory for design and inquiry rather than a validated healthcare intervention.

Conclusion

A psychologically responsive healthcare system is proposed here as one that can recognize relevant psychological and contextual conditions, interpret them without stereotyping, mobilize appropriate relational and organizational responses, support consequential participation, and revise its assumptions as circumstances

change. The framework connects behavioral science, emotional and relational care, equity, human-centered design, workforce conditions, digital accessibility, and learning while keeping their analytical levels distinct. Its central proposition is not that every patient preference should be accommodated, but that healthcare demands and system responses should be examined for usable fit within clinical, ethical, organizational, and resource boundaries. Existing evidence supports the framework's constituent premises but not the effectiveness of the integrated architecture itself. Its value therefore depends on future measurement, comparative implementation, longitudinal testing, and willingness to revise the theory when predicted relationships fail.

Acknowledgments: None

Conflict of Interest: None

Financial Support: None

Ethics Statement: None

References

1. Langberg EM, Dyhr L, Davidsen AS. Development of the concept of patient-centredness - A systematic review. *Patient Educ Couns.* 2019;102(7):1228-36. doi:10.1016/j.pec.2019.02.023
2. Bruch JD, Khazen M, Mahmic-Kaknjo M, Légaré F, Ellen ME. The effects of shared decision making on health outcomes, health care quality, cost, and consultation time: An umbrella review. *Patient Educ Couns.* 2024;129:108408. doi:10.1016/j.pec.2024.108408
3. Bombard Y, Baker GR, Orlando E, Fancott C, Bhatia P, Casalino S, et al. Engaging patients to improve quality of care: A systematic review. *Implement Sci.* 2018;13(1):98. doi:10.1186/s13012-018-0784-z
4. Boivin A, L'Espérance A, Gauvin FP, Dumez V, Macaulay AC, Lehoux P, et al. Patient and public engagement in research and health system decision making: A systematic review of evaluation tools. *Health Expect.* 2018;21(6):1075-84. doi:10.1111/hex.12804
5. Westerink HJ, Garvelink MM, van Uden-Kraan CF, Zouitni O, Bart HAJ, van der Wees PJ, et al. Evaluating patient participation in value-based healthcare: Current state and lessons learned. *Health Expect.* 2024;27(1). doi:10.1111/hex.13945
6. Panaite AC, Desroches OA, Warren É, Rouly G, Castonguay G, Boivin A. Engaging with peers to integrate community care: Knowledge synthesis and conceptual map. *Health Expect.* 2024;27(2). doi:10.1111/hex.14034
7. Cowdell F, Dyson J. How is the theoretical domains framework applied to developing health behaviour interventions? A systematic search and narrative synthesis. *BMC Public Health.* 2019;19(1):1180. doi:10.1186/s12889-019-7442-5
8. Richardson M, Khouja CL, Sutcliffe K, Thomas J. Using the theoretical domains framework and the behavioural change wheel in an overarching synthesis of systematic reviews. *BMJ Open.* 2019;9(6). doi:10.1136/bmjopen-2018-024950
9. Cabezas MF, Nazar G, Ranchor AV, Annema C. The effect of health literacy interventions on self-management in chronic diseases: A systematic review. *Ann Behav Med.* 2025;59(1). doi:10.1093/abm/kaaf073
10. Chatburn E, Marks E, Maddox L. Item development for a patient-reported measure of compassionate healthcare in action. *Health Expect.* 2024;27(1). doi:10.1111/hex.13953
11. Lerch SP, Hänggi R, Busmann Y, Lörwald A. A model of contributors to a trusting patient-physician relationship: A critical review using a systematic search strategy. *BMC Prim Care.* 2024;25(1):194. doi:10.1186/s12875-024-02435-z
12. Keshkar L, Madigan CD, Ward A, Ahmed S, Tanna V, Rahman I, et al. The effect of practitioner empathy on patient satisfaction: A systematic review of randomized trials. *Ann Intern Med.* 2024;177(2):196-209. doi:10.7326/M23-2168
13. Sinclair S, Kondejewski J, Jaggi P, Dennett L, Roze des Ordons AL, Hack TF. What is the state of compassion education? A systematic review of compassion training in health care. *Acad Med.* 2021;96(7):1057-70. doi:10.1097/ACM.0000000000004114
14. Lauwers EDL, Vandecasteele R, McMahon M, De Maesschalck S, Willems S. The patient perspective on diversity-sensitive care: A systematic review. *Int J Equity Health.* 2024;23(1):117. doi:10.1186/s12939-024-02189-1
15. Chu JN, Wong J, Bardach NS, Allen IE, Barr-Walker J, Sierra M, et al. Association between

- language discordance and unplanned hospital readmissions or emergency department revisits: A systematic review and meta-analysis. *BMJ Qual Saf.* 2024;33(7):456-69. doi:10.1136/bmjqs-2023-016295
16. Hempeler C, Schneider-Reuter L, Windel AS, Carlet J, Philipsen L, Juckel G, et al. Intersectional discrimination in mental health care: A systematic review with qualitative evidence synthesis. *Psychiatr Serv.* 2024;75(11):1125-43. doi:10.1176/appi.ps.20230252
 17. Galmarini E, Marciano L, Schulz PJ. The effectiveness of visual-based interventions on health literacy in health care: A systematic review and meta-analysis. *BMC Health Serv Res.* 2024;24(1):718. doi:10.1186/s12913-024-11138-1
 18. Wilson S, Tolley C, Mc Ardle R, Lawson L, Beswick E, Hassan N, et al. Recommendations to advance digital health equity: A systematic review of qualitative studies. *NPJ Digit Med.* 2024;7(1):173. doi:10.1038/s41746-024-01177-7
 19. Peters S, Guccione L, Francis J, Best S, Tavender E, Curran J, et al. Evaluation of research co-design in health: A systematic overview of reviews and development of a framework. *Implement Sci.* 2024;19(1):63. doi:10.1186/s13012-024-01394-4
 20. Sumner J, Ng CWT, Teo KEL, Peh ALT, Lim YW. Co-designing care for multimorbidity: A systematic review. *BMC Med.* 2024;22(1):58. doi:10.1186/s12916-024-03263-9
 21. Walker SC, Baquero B, Bekemeier B, Parnes M, Arora K. Strategies for enacting health policy codesign: A scoping review and direction for research. *Implement Sci.* 2023;18(1):44. doi:10.1186/s13012-023-01295-y
 22. Knaub M, Leon S, Pokharel S, Kiryanova V, Giroux K, Khanna P, et al. Using human-centred design to codesign patient engagement tools with a patient advisory council: Successes and challenges. *Health Expect.* 2025;28(2). doi:10.1111/hex.70230
 23. Duffy A, Boroumandzad N, Lopez Sherman A, Christie G, Riadi I, Moreno S. Examining challenges to co-design digital health interventions with end users: Systematic review. *J Med Internet Res.* 2025;27. doi:10.2196/50178
 24. Panagioti M, Geraghty K, Johnson J, Zhou A, Panagopoulou E, Chew-Graham C, et al. Association between physician burnout and patient safety, professionalism, and patient satisfaction: A systematic review and meta-analysis. *JAMA Intern Med.* 2018;178(10):1317-31. doi:10.1001/jamainternmed.2018.3713
 25. Rafi'i MR, Md Hanif SA, Daud FB. Exploring the link between healthcare organizational culture and provider work satisfaction: A systematic review. *BMC Health Serv Res.* 2025;25(1):904. doi:10.1186/s12913-025-12973-6
 26. Boskma A, van der Braak K, Hooft L, Oerbekke M, Franx A, van der Laan M. Effectiveness of organization-directed interventions on healthcare professionals' well-being: A systematic review. *EClinicalMedicine.* 2025;88:103496. doi:10.1016/j.eclinm.2025.103496
 27. Ji X, Dougherty M, Lee Y, Poghosyan L, Lelutiu-Weinberger C. Organizational interventions to address primary care provider burnout: A systematic review. *Med Care Res Rev.* 2026;83(3):167-82. doi:10.1177/10775587251391520
 28. Chang JE, Lai AY, Gupta A, Nguyen AM, Berry CA, Shelley DR. Rapid transition to telehealth and the digital divide: Implications for primary care access and equity in a post-COVID era. *Milbank Q.* 2021;99(2):340-68. doi:10.1111/1468-0009.12509
 29. Brown SH, Griffith ML, Kripalani S, Horst SN. Association of health literacy and area deprivation with initiation and completion of telehealth visits in adult medicine clinics across a large health care system. *JAMA Netw Open.* 2022;5(7). doi:10.1001/jamanetworkopen.2022.23571
 30. Borges do Nascimento IJ, Abdulazeem H, Vasanthan LT, Martinez EZ, Zucoloto ML, Østengaard L, et al. Barriers and facilitators to utilizing digital health technologies by healthcare professionals. *NPJ Digit Med.* 2023;6(1):161. doi:10.1038/s41746-023-00899-4
 31. Kim KK, Backonja U. Digital health equity frameworks and key concepts: A scoping review. *J Am Med Inform Assoc.* 2025;32(5):932-44. doi:10.1093/jamia/ocaf017
 32. de Camargo Catapan S, Sazon H, Zheng S, Gallegos-Rejas V, Mendis R, Santiago PHR, et al. A systematic review of consumers' and healthcare professionals' trust in digital healthcare. *NPJ Digit Med.* 2025;8(1):115. doi:10.1038/s41746-025-01510-8
 33. Badr J, Motulsky A, Denis JL. Digital health technologies and inequalities: A scoping review of potential impacts and policy recommendations.

- Health Policy. 2024;146:105122. doi:10.1016/j.healthpol.2024.105122
34. Wakeman M, Buckman DW, El-Toukhy S. Disparities in digital health care use in 2022. *JAMA Netw Open.* 2025;8(4). doi:10.1001/jamanetworkopen.2025.5359
 35. Silberman J, Wicks P, Patel S, Sarlati S, Park S, Korolev IO, et al. Rigorous and rapid evidence assessment in digital health with the evidence DEFINED framework. *NPJ Digit Med.* 2023;6(1):101. doi:10.1038/s41746-023-00836-5
 36. McDonald PL, Foley TJ, Verheij R, Braithwaite J, Rubin J, Harwood K, et al. Data to knowledge to improvement: Creating the learning health system. *BMJ.* 2024;384. doi:10.1136/bmj-2023-076175
 37. Lane C, McCrabb S, Turon H, Bialek C, Couper L, Wilczynska M, et al. Learning health system for implementation, scale-up, and sustainment: A systematic review to consolidate guidance for improvement. *Implement Sci.* 2026;21(1):13. doi:10.1186/s13012-025-01482-z
 38. Pomey MP, Schaad B, Lasserre-Moutet A, Böhme P, Jackson M. Towards a new integrated model for taking into account the experiential knowledge of people with chronic diseases, integrating mediation, therapeutic education and partnership: The Expanded Chronic Care Patient-Professional Partnership Model. *Health Expect.* 2024;27(5). doi:10.1111/hex.70054