

Psychosocial Interventions for Chronic Illness Adaptation and Quality of Life: Mechanisms, Effectiveness, and Future Translation Pathways

Mateo Alvarez^{1*}, Sofia Herrera², Diego Cruz¹, Andres Castro³

¹Department of Psychosocial Interventions and Chronic Illness, Faculty of Medicine, National University of Colombia, Bogota, Colombia.

²Department of Quality of Life and Adaptation, Faculty of Medicine, University of Chile, Santiago, Chile.

³Department of Translation and Psychosocial Care, Faculty of Medicine, University of Antioquia, Medellin, Colombia.

*E-mail ✉ mateo.alvarez@unal.edu.co

Abstract

Psychosocial interventions are increasingly embedded in chronic illness care, yet their effects are often interpreted as if intervention label, psychological mechanism, adaptation outcome, and implementation context were interchangeable. This review examined how psychosocial interventions influence adaptation and quality of life, which mechanisms are plausibly active, how effects vary across chronic conditions and delivery formats, and what limits translation into routine care. A systematic review was conducted using a bounded PubMed-oriented search executed on 2 September 2026 across six predefined query families spanning transdiagnostic chronic illness, chronic pain, diabetes, cancer, disease-specific conditions, and digital/implementation evidence. PRISMA 2020 guided selection reporting. Because intervention, population, comparator, outcome, and follow-up heterogeneity precluded a defensible omnibus meta-analysis, findings were synthesized using structured synthesis principles. Mechanisms, delivery characteristics, patient-reported outcomes, durability, contextual modifiers, and implementation conditions were coded separately. Seventy-two records were retrieved; three duplicates were removed, 69 unique records were screened, 45 underwent source-level eligibility assessment, and 29 evidence units were retained for the effectiveness/mechanism synthesis. Evidence supported benefits for selected psychological, behavioral, and quality-of-life outcomes, but effects were uneven across conditions, comparators, delivery modes, and follow-up periods. Mechanism evidence was strongest for constructs such as psychological flexibility, acceptance, cognitive change, self-efficacy, and guided engagement, although process change did not consistently translate into durable quality-of-life or biomedical improvement. Psychosocial intervention effectiveness is better understood as a mechanism–context configuration than as a property of a named treatment or delivery technology. Translation requires alignment among patient need, active process, outcome target, delivery support, and organizational conditions.

Keywords: Chronic illness, Psychosocial intervention, Quality of life, Psychological adaptation, Self-management, Implementation

Introduction

Chronic illness adaptation is not a single psychological endpoint. It includes emotional adjustment, sustained self-management, renegotiation of identity and roles,

symptom interpretation, social participation, and the capacity to maintain a life judged worthwhile despite ongoing disease demands. Reviews of interventions initiated around chronic-disease diagnosis show substantial diversity in intervention components and active techniques, indicating that “psychosocial intervention” is an umbrella category rather than a unitary treatment class [1]. The effectiveness of coping itself is also conditional: strategies that are adaptive under one level of perceived controllability may be ineffective or counterproductive under another, and the

Access this article online

<https://smerpub.com/>

Received: 10 May 2026; Accepted: 21 August 2026

Copyright CC BY-NC-SA 4.0

How to cite this article: Alvarez M, Herrera S, Cruz D, Castro A. Psychosocial Interventions for Chronic Illness Adaptation and Quality of Life: Mechanisms, Effectiveness, and Future Translation Pathways. *Int J Soc Psychol Asp Healthc*. 2026;6(2):167-79. <https://doi.org/10.51847/SKzTCeZVAW>

underlying evidence remains methodologically heterogeneous [2].

Cross-condition evidence illustrates both promise and restraint. Health and wellness coaching can improve quality of life, self-efficacy, and depressive outcomes, but certainty is frequently reduced by risk of bias, imprecision, and heterogeneity [3]. Professionally facilitated group work likewise produces benefits for selected psychological, self-care, and quality-of-life outcomes, although findings are mixed and some effects are short lived [4]. These limitations matter clinically because psychological comorbidity among people with chronic disease is associated with greater health-service utilization and costs [5]. At the same time, health-related quality of life is socially patterned through income, education, support, and related determinants, so intervention effects cannot be interpreted independently of the resources and constraints within which adaptation occurs [6]. Digital chronic-care research adds another layer: implementation studies frequently document feasibility and acceptability while measuring service-level consequences less consistently [7].

This review therefore distinguishes intervention content from delivery format, proposed mechanism from measured outcome, short-term response from durability, and patient-level change from organizational translation. Its central aim is not to rank named therapies across diseases, but to identify when psychosocial effects are plausible, what processes may account for them, where outcomes diverge from mechanisms, and which contextual conditions limit transfer into routine chronic care.

Materials and Methods

The review was designed as a structured systematic synthesis of psychosocial intervention evidence relevant to adult chronic physical illness. Search, screening, extraction, appraisal, and synthesis decisions were prespecified. Primary randomized trials and eligible review-level syntheses were treated as distinct evidence units; they were not pooled together or interpreted as statistically exchangeable.

Protocol and reporting standard

Selection and reporting followed PRISMA 2020 principles [8]. The review was not registered prospectively, and no unexecuted database search is represented as having occurred. The methods therefore

report the actual bounded search and verification process used for this review rather than retrospectively reconstructing a broader search universe.

Review questions

The review addressed four linked questions: which psychosocial interventions improve adaptation or quality of life in chronic illness; which psychological, behavioral, relational, or engagement mechanisms are plausibly active; how effects vary by condition, population, delivery format, comparator, and follow-up; and what implementation conditions influence translation into routine chronic care. A further analytic question asked when observed mechanism change and patient-important outcome change diverge.

Eligibility criteria

Eligible evidence was published from 2017 through 2026 in peer-reviewed Q1 journals and concerned adults living with chronic physical illness. Included primary studies evaluated a psychosocial or psychological intervention and reported adaptation-relevant psychological, behavioral, functional, or quality-of-life outcomes. Eligible review-level evidence synthesized comparable intervention questions or provided directly relevant mechanism, durability, or implementation evidence. Records were excluded when they were primarily biomedical, outside the chronic-illness adaptation scope, lacked usable psychosocial outcomes, were protocol-only or preliminary without adequate outcome evidence, or failed the journal/design eligibility rules.

Information sources

Searches were executed on 2 September 2026 using PubMed-oriented retrieval. Bibliographic and DOI metadata were verified against PubMed and publisher records, with citation chaining used to confirm directly relevant trials and syntheses. Journal quartile eligibility was checked in the relevant subject category and publication year where available, using the latest available annual ranking for 2025–2026 publications when a same-year annual quartile was not yet released.

Search strategy

Six query families were used: transdiagnostic chronic illness with psychosocial intervention and adaptation/quality of life; chronic pain with ACT, mindfulness, CBT, and psychological outcomes; diabetes with distress, self-management, psychological

intervention, and quality of life; cancer with digital or psychological intervention and distress/quality of life; condition-specific searches for inflammatory bowel disease, multiple sclerosis, COPD, and multimorbidity; and digital-health searches focused on chronic-disease implementation and translation. Searches were intentionally broad enough to retrieve null, heterogeneous, and implementation-focused evidence rather than only supportive intervention studies.

Study selection

Across the six search families, 72 records were retrieved. Three duplicate appearances were removed, leaving 69 unique records. Title and abstract screening excluded 24 records, and 45 records underwent source-level eligibility assessment. Sixteen were then excluded because of journal/design ineligibility, insufficient mechanism or adaptation fit, preliminary evidence status, or redundancy with stronger eligible syntheses. Twenty-nine evidence units entered the effectiveness/mechanism synthesis. Eight additional methodological, conceptual,

contextual, overlap, or implementation references supported interpretation without being counted as effectiveness evidence.

Data extraction

For each evidence unit, extraction separated design, chronic condition, population or evidence unit, intervention content, delivery format, comparator, proposed mechanism, measurement approach, psychological and quality-of-life outcomes, behavioral or functional outcomes, follow-up, contextual modifiers, risk-of-bias information, and implementation implications. Missing information was retained as unreported rather than inferred. Delivery technology was not coded as a mechanism, and process measures were not treated as clinical outcomes.

Because apparent intervention heterogeneity can arise from different sources, the extraction framework explicitly preserves the distinctions summarized in **Table 1**.

Table 1. Heterogeneity map for interpreting psychosocial intervention evidence across chronic illness

Extraction condition	What is held separate	Why the distinction matters	Consequence for synthesis	Main uncertainty
Patient entry state	Diagnosis stage, baseline distress, symptom burden, multimorbidity	Identical interventions may address different adaptation problems	Compare effects within clinically similar entry states	Regression to mean and self-selection
Intervention architecture	Therapy content, support intensity, guidance, session structure	A digital label can conceal different active components	Classify content before delivery channel	Multicomponent attribution
Comparator strength	Waitlist, usual care, attention control, active psychotherapy	Comparative effect depends on what the intervention is tested against	Give active-comparator findings greater comparative weight	Expectancy and contact effects
Mechanism measurement	Flexibility, acceptance, self-efficacy, cognitive change, emotional processing	Process change is not equivalent to outcome change	Report mechanism and endpoint evidence separately	Mediation rarely established
Outcome layer	Distress, behavior, function, quality of life, biomedical status	Improvement can occur at one layer without transferring to another	Avoid composite judgments of “success”	Instrument and timing differences
Temporal window	Immediate, short-term, longer follow-up	Adaptation effects may attenuate or consolidate	Interpret durability independently from initial response	Inconsistent follow-up intervals
Care context	Specialty clinic, primary care, digital self-management, community delivery	Workflow and access can enable or constrain effect	Separate efficacy from translation conditions	System-specific generalizability

Intervention taxonomy

Interventions were coded first by therapeutic content and only then by delivery format. The resulting taxonomy

included cognitive-behavioral approaches, acceptance- and mindfulness-based approaches, emotionally focused interventions, coaching and self-management support,

facilitated group interventions, and multicomponent digital programs. Delivery was coded separately as face-to-face, group-based, telephone, web/app-based, guided digital, unguided digital, or hybrid. This prevents the analytically weak assumption that two interventions delivered through the same technology share the same active mechanism.

Mechanism-of-action coding

Mechanisms were coded when a source explicitly measured or theoretically specified a process relevant to adaptation. Candidate processes included psychological flexibility, acceptance, cognitive restructuring, emotional awareness and expression, self-efficacy, coping fit, engagement, and relational support. Evidence was classified as stronger when process change was measured temporally and linked to patient-important outcomes, and weaker when a mechanism was inferred only from treatment branding. No process was treated as a validated mediator unless the source directly supported mediation.

Risk-of-bias assessment

Source-reported risk-of-bias, certainty, and review-quality assessments were retained. Interpretation additionally distinguished inactive from active comparators, short from longer follow-up, self-report from functional or clinical outcomes, and primary trials from secondary syntheses. These distinctions were used as interpretive flags rather than converted into a new numerical quality score.

Effectiveness synthesis

Because the evidence spans different diseases, intervention families, comparators, outcomes, and follow-up periods, an omnibus meta-analysis was not considered scientifically defensible. Structured synthesis followed SWiM principles [9], grouping evidence by intervention architecture, mechanism, outcome layer, disease context, comparator, and durability. Quantitative findings reported by eligible reviews were retained as source-level evidence, but no new pooled effect was calculated across non-equivalent evidence units.

Results and Discussion

The 29 synthesis-eligible evidence units encompassed primary randomized evidence and review-level syntheses across transdiagnostic chronic illness, chronic pain, diabetes, cancer, inflammatory bowel disease, multiple sclerosis, COPD, and multimorbidity. The evidence base was sufficiently diverse to evaluate psychological adaptation, self-management, quality of life, delivery effects, durability, and translation, but it was not sufficiently uniform to support a single estimate of psychosocial intervention effectiveness.

The selection logic is shown in **Figure 1** because the distinction between synthesis-eligible evidence and contextual or methodological support is essential to interpreting the review denominator.




Figure 1. Review-selection logic separating synthesis evidence from contextual support

Study selection

The 24 title/abstract exclusions comprised records outside the publication window, primarily biomedical or nonpsychosocial studies, records without a relevant chronic-illness adaptation construct, and records lacking an eligible intervention/outcome or preliminary

journal/design fit. At source-level assessment, exclusions reflected failure of the Q1/design rule, insufficient fit to psychosocial mechanism or adaptation outcomes, protocol/pilot evidence without adequate outcomes, or redundancy with stronger eligible syntheses. The consequences of these selection decisions for interpretation are summarized in **Table 2**.

Table 2. Measurement consequences of evidence-unit selection in the chronic-illness psychosocial synthesis

Evidence pattern admitted to synthesis	What it can contribute	Measurement consequence	Interpretive boundary	Use in this review
Randomized trial with active comparator	Comparative psychosocial effect under matched contact	Isolates treatment-specific difference more strongly	Cannot establish broad generalizability	Comparative effectiveness
Randomized trial with inactive/waitlist comparator	Efficacy signal against minimal intervention	Susceptible to expectancy/contact differences	Superiority over another credible therapy is not established	Initial effectiveness
Systematic review/meta-analysis	Cross-study direction and heterogeneity	Inherits variation in measures and trial quality	Pooled estimate may conceal context	Outcome synthesis
Network meta-analysis	Indirect and direct comparative structure	Rankings depend on network assumptions	Ranking is not universal clinical superiority	Modality comparison
Umbrella/overview evidence	Review-level consistency and overlap	Identifies redundancy and certainty problems	Does not add independent participants or trials	Evidence-boundary analysis
Implementation/qualitative evidence	Workflow, acceptability, policy, and organizational conditions	Captures translation domains poorly measured in efficacy trials	Cannot estimate treatment effect	Routine-care translation

Study characteristics

Evidence units varied not only by chronic condition but by adaptation state. Some addressed newly diagnosed or broadly defined chronic disease, whereas others enrolled people selected for distress, insomnia, pain, disability, or disease-specific psychosocial burden. This heterogeneity means that apparently different treatment effects may partly reflect different entry problems rather than different intervention potency. Digital and remotely

delivered interventions were represented across both transdiagnostic and disease-specific contexts, and review-level evidence ranged from focused meta-analysis to umbrella synthesis, making evidence overlap itself an interpretive issue.

Table 3 converts these differences into clinically meaningful context states rather than presenting a generic study inventory.

Table 3. Chronic-illness adaptation states that alter interpretation of psychosocial intervention effects

Dominant adaptation problem	Mechanism most plausibly targeted	Outcome that best tests benefit	Competing explanation or uncertainty	Translation implication
Care-context state — Early post-diagnosis adaptation				
Uncertainty, role disruption, coping reorganization	Coping fit, self-efficacy, meaning-making	Distress, adjustment, participation	Spontaneous adaptation over time	Timing and readiness matter
Care-context state — High psychological distress				

Anxiety, depression, illness-specific distress	Cognitive change, acceptance, emotion regulation	Symptom burden and mental QoL	Nonspecific support/contact effects	Screening can improve targeting
Care-context state — Persistent pain/disability				
Interference, avoidance, restricted valued activity	Flexibility, acceptance, cognitive/emotional processing	Function, interference, life participation	Pain intensity may not change in parallel	Outcome choice must match mechanism
Care-context state — Self-management burden				
Sustained daily decisions and treatment work	Self-efficacy, problem solving, engagement	Behavior, distress, treatment burden	Biomedical markers may remain unchanged	Integrate with clinical management
Care-context state — Survivorship/long-term treatment				
Recurrence fear, sleep disturbance, cumulative burden	Coping, CBT, mindfulness, guided support	QoL, sleep, distress	Improvements may attenuate	Maintenance strategy needed
Care-context state — Multimorbidity or fragmented care				
Competing conditions and treatment priorities	Prioritization, agency, coordinated support	Function, QoL, care burden	Single-condition programs may misfit	Organizational coordination forms part of intervention context

Intervention types and delivery formats

Delivery modality did not map cleanly onto therapeutic mechanism. A waitlist-controlled digital ACT trial in adults with chronic health conditions found early improvement in psychological flexibility and later differences in quality of life and impairment, while therapist-supported digital CBT and telephone-delivered CBT likewise showed psychological benefits in chronic-condition populations; these findings support remote delivery as feasible but do not imply mechanism equivalence [10-12]. Connected-health cancer evidence is especially heterogeneous, with many interventions but only a minority of trials sufficiently comparable for pooled analysis [13]. More recent digital cancer syntheses show benefits for selected distress and quality-of-life outcomes while also indicating that modality and

human guidance alter engagement and efficacy [14-16]. An umbrella review further shows that the apparent breadth of digital benefit sits within a review literature of variable methodological quality [17]. Self-help interventions in diabetes demonstrate the same need to separate delivery from outcome timing: psychological, behavioral, and quality-of-life benefits vary by follow-up interval [18]. Earlier broad psychological-intervention evidence did not establish firm effects on diabetes-related distress or health-related quality of life [19], whereas more recent distress-targeted synthesis indicates stronger short-term distress improvement without consistent long-term distress or glycemic benefit [20]. **Table 4** therefore crosswalks intervention architecture against the interpretation it can sustain.

Table 4. Evidence–interpretation crosswalk for psychosocial intervention architecture and delivery

Intervention architecture	Delivery forms represented	Primary process target	Outcome signal most defensible	Context modifier	Interpretation that should be avoided
Cognitive-behavioral	Therapist-supported digital, telephone, group, telehealth	Appraisal and behavioral response	Distress, coping, function	Therapist contact and disease specificity	Treating all CBT delivery as equivalent
Acceptance/mindfulness based	Digital, hybrid, group, individual	Flexibility, acceptance, present-moment regulation	Distress, function, mental QoL	Baseline distress and practice engagement	Assuming process change guarantees clinical change
Guided digital support	Web/app with human guidance	Engagement plus therapy-specific process	Selected distress/QoL outcomes	Guidance intensity and access	Attributing benefit to technology alone
Self-help/coaching	Structured self-management, remote or mixed support	Self-efficacy and problem solving	Behavior, distress, QoL	Readiness and treatment burden	Expecting uniform biomedical improvement

Facilitated group intervention	Professionally led group sessions	Shared learning, normalization, support	Psychological and self-care outcomes	Group composition and facilitation	Assuming social contact itself is the active ingredient
Condition-specific psychosocial care	Specialty-tailored CBT/ACT/mindfulness	Disease-relevant distress or disability process	Illness-specific function and QoL	Symptom phase and specialty workflow	Generalizing directly across diseases

Theoretical foundations

The reviewed interventions draw on partially overlapping but analytically distinct theories. Coping-fit models emphasize matching coping strategy to perceived controllability, whereas CBT-oriented approaches target interpretations and behavioral responses. ACT focuses more explicitly on psychological flexibility and values-consistent action in the presence of difficult internal experiences. Earlier chronic-pain synthesis found that ACT can improve acceptance, flexibility, functioning, and mood even when quality of life or pain intensity does not improve consistently [21]. Later review work reinforced psychological flexibility as a plausible process while showing that moderator and predictor evidence remains limited [22]. A three-level meta-analysis further suggested that ACT effects vary with delivery mode, diagnosis, and follow-up duration [23], while an overview of systematic reviews identified substantial overlap and uneven certainty across the ACT evidence base [24].

Process evidence therefore cannot be collapsed into a single “flexibility” construct. A 2026 synthesis found improvements in global psychological flexibility and acceptance, but not consistent improvement in committed action or values, with low or very low certainty for several outcomes [25]. Cognitive restructuring provides a contrasting mechanism: a meta-analysis in chronic pain reported favorable pain effects but also substantial methodological variability [26]. Taken together, these theories support a process-based interpretation in which intervention names are treated as imperfect proxies for active psychological mechanisms. The review therefore carries forward a proposed distinction among mechanism engagement, patient-reported adaptation, behavioral transfer, quality-of-life change, and clinical consequence; subsequent sections test where those layers converge and where they separate.

Active psychosocial mechanisms

Across the evidence base, psychosocial mechanisms were more consistently demonstrated at the level of

patient-reported process than at the level of downstream clinical consequence. Psychological flexibility and acceptance were the most explicitly measured processes within ACT-oriented studies. However, the construct was not uniform: global flexibility and acceptance improved more consistently than committed action or values-based behavior, and certainty around several process outcomes remained low [25]. Cognitive restructuring represents a different pathway, directed toward appraisal and interpretation rather than willingness to experience difficult internal states; chronic-pain synthesis suggests potential benefit, although study quality and intervention composition varied substantially [26].

Other mechanisms were less directly tested. Self-efficacy plausibly links coaching or self-management support to behavior, whereas guidance may increase engagement with digital interventions. Emotional awareness and expression may alter avoidance or affective processing, and professionally facilitated groups may provide normalization and interpersonal reinforcement. Yet these mechanisms cannot be assumed from treatment labels alone. The review therefore distinguishes mechanism engagement from mechanism validation: the former requires evidence that a targeted psychological process changed, whereas the latter would require stronger temporal or mediation evidence demonstrating that the process accounts for subsequent adaptation or quality-of-life improvement.

Emotional and psychological adaptation outcomes

Psychological outcomes were generally more responsive than biomedical outcomes, but response depended strongly on comparator and population. In older veterans with musculoskeletal chronic pain, Emotional Awareness and Expression Therapy produced greater reductions in pain severity and improvements in several secondary psychological outcomes than an active CBT comparator, suggesting that emotionally focused mechanisms may be clinically relevant in selected populations [27]. Such evidence is comparatively

informative because it tests one credible psychosocial treatment against another rather than against minimal care.

Null findings were equally important. Among cancer survivors with insomnia, a behavioral sleep intervention did not produce significant between-group advantages in health-related quality of life, depression, anxiety, or mood relative to an attention control despite within-group improvement [28]. This illustrates why change over time cannot automatically be attributed to the intervention and why psychological adaptation should not be inferred from uncontrolled pre–post change.

Social and relational outcomes

Relational outcomes were measured less frequently than distress or quality of life. Group interventions and guided digital programs suggest that professional support, shared experience, normalization, and interpersonal accountability may facilitate engagement, but these relational conditions are rarely isolated experimentally. The distinction matters because increased therapist contact can improve adherence or perceived support without necessarily representing the therapy-specific mechanism.

Accordingly, the current evidence supports relational support primarily as an enabling condition rather than a universally established mediator. Future studies need to distinguish whether human guidance changes exposure to the intervention, strengthens therapeutic alliance, modifies self-efficacy, or independently affects adaptation.

Self-management and behavioral outcomes

Behavioral change was most evident when interventions explicitly addressed self-management, coping, or daily disease work. Coaching and structured self-help can improve self-efficacy or health behavior, but psychological improvement did not reliably produce parallel change in physiological endpoints. Diabetes evidence was especially illustrative: improvements in distress or self-management measures could occur without sustained glycemic improvement.

This separation argues against evaluating psychosocial interventions solely through biomarkers. A psychosocial intervention may be successful if it reduces distress, improves functional participation, or makes treatment demands more manageable even when the underlying disease marker remains unchanged. Conversely, a

biomedical improvement cannot by itself establish that the proposed psychosocial mechanism was active.

Quality-of-life outcomes

Quality of life was the most integrative but also one of the most difficult outcomes to interpret because instruments combine psychological, functional, social, and sometimes symptom-related domains. In multiple sclerosis, mindfulness-based cognitive therapy and cognitive rehabilitation improved selected mental quality-of-life and depressive outcomes, but anxiety, physical quality of life, and everyday cognitive functioning were less consistently affected, and several benefits attenuated over follow-up [29].

Across the wider evidence base, quality-of-life effects were therefore best regarded as outcome-specific and context-dependent. Improvements were observed in selected ACT, CBT, coaching, cancer-support, and disease-specific interventions, yet null or modest findings occurred in other studies. The review found no basis for treating quality-of-life improvement as an automatic consequence of reducing psychological distress.

Differential effects across conditions and populations

Condition-specific evidence reinforced the importance of context. Recent inflammatory bowel disease trials reported psychosocial benefits across group CBT, mindfulness-based cognitive therapy, and ACT, but clinical consequences differed: psychological functioning and quality of life could improve without corresponding reductions in flare occurrence or without superiority across all secondary outcomes [30–32]. Telehealth IBD-specific CBT similarly reduced disability while leaving disease activity and costs comparatively unchanged [33].

In stable COPD, network meta-analysis suggested advantages for some nonpharmacological approaches for anxiety and depression, but certainty was low and publication bias remained possible [34]. Across conditions, baseline distress, disability, age, disease activity, multimorbidity, treatment burden, digital access, guidance intensity, and clinical setting therefore function as plausible effect modifiers. These variables should not be interpreted as fixed patient traits; they define circumstances under which intervention demands and available resources may fit more or less well.

Durability of effects

Evidence for sustained benefit was thinner than evidence for initial response. Group interventions, diabetes-distress treatment, ACT, cancer survivorship programs, and multiple-sclerosis interventions all showed examples in which short-term gains diminished, became nonsignificant, or remained outcome-specific over longer follow-up. Durability therefore represents a separate effectiveness question rather than an extension of immediate efficacy.

A useful distinction is between initial adaptation and maintenance of adaptation. Initial change may depend on therapeutic learning or emotional relief, whereas maintenance may require repeated practice, environmental reinforcement, ongoing clinical support, or modification of the intervention as illness circumstances change.

Implementation and translation evidence

Translation into routine care requires more than demonstrating efficacy. Stakeholder evidence on person-centered planning for patients with multiple chronic conditions identifies workflow redesign, interdisciplinary coordination, payment structures, information systems, and organizational support as conditions affecting implementation [35]. Digital multimorbidity interventions likewise show inconsistent physiological, psychological, and functional benefits, with low or very low certainty for several outcomes and substantial variation between disease combinations and settings [36]. European primary-care evidence similarly found little advantage of digital interventions for quality of life, depression, anxiety, and several other outcomes [37].

These findings indicate that technology should be treated as an access and delivery infrastructure rather than an active psychosocial mechanism. Translation depends on whether intervention demands fit patient capability, whether clinicians can act on generated information, whether support is available when engagement falls, and whether the intervention integrates with existing chronic-care work.

The reviewed literature supports psychosocial intervention as an important component of chronic illness care, but it does not support a single transdiagnostic estimate of effectiveness. Effects differ according to therapeutic process, comparator, outcome layer, disease context, delivery support, and follow-up. Psychological outcomes often improve more readily than clinical disease measures, while quality-of-life effects occupy an intermediate position because they integrate emotional, functional, and social consequences.

The central synthesis is therefore that psychosocial interventions should be understood as mechanism–context configurations. A named therapy is not itself a mechanism, a digital platform is not itself an intervention mechanism, and improvement in one outcome does not prove transfer across outcome layers. This interpretation accommodates both positive and null evidence without treating heterogeneous results as methodological noise alone.

Mechanisms versus outcomes

The most important distinction emerging from the review is between process activation and consequential adaptation. Psychological flexibility, acceptance, cognitive change, emotional processing, engagement, or self-efficacy may improve without producing durable quality-of-life or biomedical effects. Conversely, a patient-important outcome may improve even when the hypothesized mediator was not measured or did not change clearly.

Mechanism–outcome discordance appears repeatedly in the evidence: distress can improve without glycemic change; psychosocial wellbeing can improve without reduced inflammatory activity; disability can improve without measurable disease modification; and quality-of-life improvement may attenuate over time. Mechanisms should therefore be interpreted within nested levels extending from psychological process through behavior and lived functioning to clinical and organizational consequence.

The scale-transfer boundaries underlying this interpretation are summarized in **Figure 2**.




Figure 2. Where psychosocial change stops transferring: nested boundaries between mechanism engagement and consequential adaptation

Translation into routine chronic care

Routine implementation should begin with the clinical problem rather than the technology or therapy label. A patient whose principal problem is distress may require a different mechanism and outcome target from one whose principal difficulty is avoidance, treatment burden, disability, or fragmented self-management. Delivery should then be matched to access, preference, digital capability, available professional support, and the capacity of the clinical service to respond.

This approach favors adaptive psychosocial care over uniform program deployment. Guidance may be intensified when engagement falls; intervention targets may shift as illness circumstances change; and outcome monitoring should distinguish psychological, behavioral, functional, and clinical response rather than relying on one composite judgment of success.

Strengths and limitations

Strengths include a deliberately transdiagnostic scope, preservation of active-comparator and null evidence, explicit separation of mechanisms from outcomes, and inclusion of implementation and durability evidence. The

review also avoids combining primary trials and secondary syntheses into a single quantitative estimate. Limitations are substantial. The search was a bounded PubMed-oriented retrieval rather than an exhaustive multi-database export. Evidence density differed markedly by condition, several topics were represented predominantly by secondary syntheses, and overlap among reviews limits apparent independence. Intervention terminology and outcome measurement were inconsistent, follow-up was often short, and process measures were commonly self-reported. Mechanistic mediation was rarely established, and underserved populations, resource-limited settings, language barriers, accessibility constraints, and structural inequities were incompletely represented.

Future research

Future trials should specify the mechanism expected to change, measure that process before downstream outcomes, and test whether process change mediates adaptation rather than assuming mechanism from intervention identity. Active comparators, longer follow-

up, standardized patient-important outcomes, and repeated mechanism measurement are especially needed. Research should also test adaptive treatment matching: whether intervention content, guidance intensity, and delivery format can be altered according to baseline distress, multimorbidity, treatment burden, social resources, or early response. Implementation studies should measure workflow consequences, clinician uptake, equity, accessibility, costs, and sustained service integration alongside patient outcomes. Such work would permit the proposed mechanism–context architecture to move from explanatory synthesis toward prospective validation.

Conclusion

Psychosocial interventions can improve important dimensions of living with chronic illness, but their effectiveness is neither uniform nor reducible to therapy name or delivery technology. The evidence is strongest when the intervention target, psychological mechanism, patient-important outcome, and clinical context are closely aligned. Psychological processes may improve without durable quality-of-life or biomedical transfer, and favorable outcomes do not by themselves validate a proposed mechanism. Chronic illness psychosocial care should therefore be evaluated as a conditional, multilevel process linking mechanism engagement to adaptation through behavioral, relational, social, clinical, and organizational boundaries. Future translation depends on testing these boundaries directly and designing interventions that can adapt as patient needs and illness contexts change.

Acknowledgments: None

Conflict of Interest: None

Financial Support: None

Ethics Statement: None

References

1. Akyirem S, Forbes A, Wad JL, Due-Christensen M. Psychosocial interventions for adults with newly diagnosed chronic disease: A systematic review. *J Health Psychol.* 2022;27(7):1753-82. doi:10.1177/1359105321995916
2. Finkelstein-Fox L, Park CL. Control-coping goodness-of-fit and chronic illness: A systematic review of the literature. *Health Psychol Rev.* 2019;13(2):137-62. doi:10.1080/17437199.2018.1560229
3. Boehmer KR, Álvarez-Villalobos NA, Barakat S, de Leon-Gutierrez H, Ruiz-Hernandez FG, Elizondo-Omaña GG, et al. The impact of health and wellness coaching on patient-important outcomes in chronic illness care: A systematic review and meta-analysis. *Patient Educ Couns.* 2023;117:107975. doi:10.1016/j.pec.2023.107975
4. Jackson M, Jones D, Dyson J, Macleod U. Facilitated group work for people with long-term conditions: A systematic review of benefits from studies of group-work interventions. *Br J Gen Pract.* 2019;69(682). doi:10.3399/bjgp19X702233
5. Sporinova B, Manns B, Tonelli M, Hemmelgarn B, MacMaster F, Mitchell N, et al. Association of mental health disorders with health care utilization and costs among adults with chronic disease. *JAMA Netw Open.* 2019;2(8). doi:10.1001/jamanetworkopen.2019.9910
6. Kangas T, Milis SL, Vanthomme K, Vandenheede H. The social determinants of health-related quality of life among people with chronic disease: A systematic literature review. *Qual Life Res.* 2025;34(9):2501-11. doi:10.1007/s11136-025-03976-1
7. Pong C, Tseng RMWW, Tham YC, Lum E. Current implementation of digital health in chronic disease management: Scoping review. *J Med Internet Res.* 2024;26. doi:10.2196/53576
8. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: An updated guideline for reporting systematic reviews. *BMJ.* 2021;372. doi:10.1136/bmj.n71
9. Campbell M, McKenzie JE, Sowden A, Katikireddi SV, Brennan SE, Ellis S, et al. Synthesis without meta-analysis (SWiM) in systematic reviews: Reporting guideline. *BMJ.* 2020;368. doi:10.1136/bmj.l6890
10. Levin ME, Aller TB, Klimczak KS, Donahue ML, Knudsen FM. Digital acceptance and commitment therapy for adults with chronic health conditions: Results from a waitlist-controlled trial. *Behav Res Ther.* 2025;188:104729. doi:10.1016/j.brat.2025.104729

11. Picariello F, Hulme K, Seaton N, Hudson JL, Norton S, Wroe A, et al. A randomized controlled trial of a digital cognitive-behavioral therapy program (COMPASS) for managing depression and anxiety related to living with a long-term physical health condition. *Psychol Med.* 2024;54(8):1796-809. doi:10.1017/S0033291723003756
12. Altieri M, Sergi MR, Tommasi M, Santangelo G, Saggino A. The efficacy of telephone-delivered cognitive behavioral therapy in people with chronic illnesses and mental diseases: A meta-analysis. *J Clin Psychol.* 2024;80(1):223-54. doi:10.1002/jclp.23563
13. Gitonga I, Desmond D, Duda N, Maguire R. Impact of connected health interventions on psychological wellbeing and quality of life in patients with cancer: A systematic review and meta-analysis. *Psychooncology.* 2022;31(10):1621-36. doi:10.1002/pon.6019
14. Zhang T, Ren Z, Wakefield CE, Hui BPH, Akechi T, Shi C, et al. Are digital psychological interventions for psychological distress and quality of life in cancer patients effective? A systematic review and network meta-analysis. *Clin Psychol Rev.* 2025;115:102520. doi:10.1016/j.cpr.2024.102520
15. Akdemir A, Smith AB, Wu VS, Rincones O, Russell H, Lyhne JD, et al. Guided versus non-guided digital psychological interventions for cancer patients: A systematic review and meta-analysis of engagement and efficacy. *Psychooncology.* 2024;33(1). doi:10.1002/pon.6290
16. Dong M, Huang Y, Zhang X, Wu L, Ghazal L, Cao F. Effects of psychological intervention based on digital health technology on psychological outcomes in cancer survivors: Systematic review and meta-analysis. *Psychooncology.* 2025;34(5). doi:10.1002/pon.70166
17. Zhong C, Luo X, Tan M, Chi J, Guo B, Tang J, et al. Digital health interventions to improve mental health in patients with cancer: Umbrella review. *J Med Internet Res.* 2025;27. doi:10.2196/69621
18. Wicaksana AL, Apriliyasari RW, Tsai PS. Effect of self-help interventions on psychological, glycemic, and behavioral outcomes in patients with diabetes: A meta-analysis of randomized controlled trials. *Int J Nurs Stud.* 2024;149:104626. doi:10.1016/j.ijnurstu.2023.104626
19. Chew BH, Vos RC, Metzendorf MI, Scholten RJPM, Rutten GEHM. Psychological interventions for diabetes-related distress in adults with type 2 diabetes mellitus. *Cochrane Database Syst Rev.* 2017;9(9). doi:10.1002/14651858.CD011469.pub2
20. Zu W, Zhang S, Du L, Huang X, Nie W, Wang L. The effectiveness of psychological interventions on diabetes distress and glycemic level in adults with type 2 diabetes: A systematic review and meta-analysis. *BMC Psychiatry.* 2024;24(1):660. doi:10.1186/s12888-024-06125-z
21. Hughes LS, Clark J, Colclough JA, Dale E, McMillan D. Acceptance and commitment therapy (ACT) for chronic pain: A systematic review and meta-analyses. *Clin J Pain.* 2017;33(6):552-68. doi:10.1097/AJP.0000000000000425
22. McCracken LM, Yu L, Vowles KE. New generation psychological treatments in chronic pain. *BMJ.* 2022;376. doi:10.1136/bmj-2021-057212
23. Lai L, Liu Y, McCracken LM, Li Y, Ren Z. The efficacy of acceptance and commitment therapy for chronic pain: A three-level meta-analysis and a trial sequential analysis of randomized controlled trials. *Behav Res Ther.* 2023;165:104308. doi:10.1016/j.brat.2023.104308
24. Martinez-Calderon J, García-Muñoz C, Rufo-Barbero C, Matias-Soto J, Cano-García FJ. Acceptance and commitment therapy for chronic pain: An overview of systematic reviews with meta-analysis of randomized clinical trials. *J Pain.* 2024;25(3):595-617. doi:10.1016/j.jpain.2023.09.013
25. Sanabria-Mazo JP, Rodríguez-Freire C, Royuela-Colomer E, Ponce-López P, Alonso-Coello P, Barrios V, et al. Do acceptance- and mindfulness-based interventions improve psychological flexibility in people with chronic pain? A systematic review and meta-analysis of randomized controlled trials. *Eur J Pain.* 2026;30(7). doi:10.1002/ejp.70342
26. Pinteá S, Maier P. Mind over chronic pain: A meta-analysis of cognitive restructuring in chronically ill adults. *J Psychosom Res.* 2024;184:111837. doi:10.1016/j.jpsychores.2024.111837
27. Yarns BC, Jackson NJ, Alas A, Melrose RJ, Lumley MA, Sultzer DL. Emotional awareness and expression therapy vs cognitive behavioral therapy for chronic pain in older veterans: A randomized clinical trial. *JAMA Netw Open.* 2024;7(6). doi:10.1001/jamanetworkopen.2024.15842
28. Kwon M, Zhu J, Wilding GE, Larkin K, Gehrman PR, Dickerson SS. Health-related quality of life and

- mental health outcomes among cancer survivors in an insomnia intervention: A randomized controlled trial. *Ann Behav Med.* 2025;59(1). doi:10.1093/abm/kaae096
29. Nauta IM, van Dam M, Bertens D, Kessels RPC, Fasotti L, Uitdehaag BMJ, et al. Improved quality of life and psychological symptoms following mindfulness and cognitive rehabilitation in multiple sclerosis and their mediating role for cognition: A randomized controlled trial. *J Neurol.* 2024;271(7):4361-72. doi:10.1007/s00415-024-12327-y
 30. Kalogeropoulou M, Karaivazoglou K, Konstantopoulou G, Vinni E, Sotiropoulos C, Tourkochristou E, et al. The impact of group cognitive behavioral psychotherapy on disease severity and psychosocial functioning in patients with inflammatory bowel disease: A randomized controlled study. *J Crohns Colitis.* 2025;19(3). doi:10.1093/ecco-jcc/jjae144
 31. Ter Avest MM, Huijbers MJ, Horjus CS, Römkens TEH, Witterman EM, van Dop WA, et al. Group-delivered mindfulness-based cognitive therapy to reduce psychological distress and improve sleep in patients with inflammatory bowel diseases: A multicenter randomized controlled trial (MindIBD). *Inflamm Bowel Dis.* 2025;31(11):3021-32. doi:10.1093/ibd/izaf116
 32. Naude C, Skvarc D, Maunick B, Evans S, Romano D, Chesterman S, et al. Acceptance and commitment therapy for adults living with inflammatory bowel disease and distress: A randomized controlled trial. *Am J Gastroenterol.* 2025;120(8):1839-51. doi:10.14309/ajg.0000000000003032
 33. Tse CS, Lewis JD, Horst SN, Saha S, Taleban S, Shah SA, et al. Inflammatory bowel diseases-specific cognitive behavioral therapy delivered through telehealth reduces disability: Addressing Disability Effectively With Psychosocial Telemedicine pragmatic randomized controlled trial. *Am J Gastroenterol.* 2026;121(6):1435-44. doi:10.14309/ajg.0000000000003681
 34. Xue Y, Tang K, Liu H, Jin X. Efficacy of non-pharmacological interventions on psychological well-being in stable COPD: A systematic review and network meta-analysis. *J Psychosom Res.* 2025;196:112324. doi:10.1016/j.jpsychores.2025.112324
 35. Watson BN, Estenson L, Eden AR, Gerstein MT, Carney MT, Dotson VM, et al. Person-centered care planning for people living with or at risk for multiple chronic conditions. *JAMA Netw Open.* 2024;7(10). doi:10.1001/jamanetworkopen.2024.39851
 36. Qiu R, Feng H, Li X, Zhou H. Effectiveness and implementation of digital health interventions on physiological, psychological, and functional outcomes in adults with multimorbidity: Systematic review and meta-analysis of randomized controlled trials. *J Med Internet Res.* 2026;28. doi:10.2196/90458
 37. Ambrosi E, Mezzalana E, Canzan F, Leardini C, Vita G, Marini G, et al. Effectiveness of digital health interventions for chronic conditions management in European primary care settings: Systematic review and meta-analysis. *Int J Med Inform.* 2025;196:105820. doi:10.1016/j.ijmedinf.2025.105820